

DISCRETE INTERPOLATION BETWEEN MONOTONE PROBABILITY AND FREE PROBABILITY¹

Romuald Lenczewski and Rafał Śaławata

*Institute of Mathematics
Wrocław University of Technology
Wybrzeże Wyspiańskiego 27
50-370 Wrocław
Poland*

Abstract

We construct a sequence of states called *m-monotone product states* which give a discrete interpolation between the monotone product of states of Muraki and the free product of states of Avitzour and Voiculescu in free probability. We derive the associated basic limit theorems and develop the combinatorics based on non-crossing ordered partitions with monotone order starting from depth m . The Hilbert space representations of the limit mixed moments in the invariance principle lead to *m-monotone Gaussian operators* living in *m-monotone Fock spaces*, which are truncations of the free Fock space $\mathcal{F}(L^2(\mathbb{R}^+))$. A new type of combinatorics of inner blocks in non-crossing partitions leads to explicit formulas for the mixed moments of *m-monotone Gaussian operators*, which are new even in the case of monotone independent Gaussian operators with arcsine distributions.

1. INTRODUCTION.

The aim of this paper is to construct an interpolation between free probability [V,VDN] based on the notion of the free product of states [Av,V] and monotone probability based on the notion of the monotone product of states [M2]. Our interpolation is given by a sequence of product states called *m-monotone*, where $m \in \mathbb{N}$, which gives the monotone product for $m = 1$, whose asymptotics leads to the free product for $m = \infty$.

The definition of the free product of states is based on the freeness condition

$$\phi(a_1 a_2 \dots a_n) = 0 \quad \text{for } a_k \in \mathcal{A}_{i_k} \cap \text{Ker} \phi_{i_k} \quad (1.1)$$

on the free product $\ast_{i \in I} \mathcal{A}_i$ of unital $\ast$ -algebras with identified units ($1_i = 1_k$ for all $i, k \in I$), where $i_1 \neq i_2 \neq \dots \neq i_n$.

¹This work is supported by KBN grant No 2P03A00723 and by the EU Network QP-Applications, Contract No. HPRN-CT-2002-00279

It is interesting to observe that the monotone product of states can also be defined using (1.1), except that one has to take the free product without identification of units $\sqcup_{i \in I} \mathcal{A}_i$, provided I is linearly ordered, and give conditions on moments involving units, namely

$$\phi(a_1 \dots a_{k-1} 1_{i_k} a_{k+1} \dots a_n) = \phi(a_1 \dots a_{k-1} a_{k+1} \dots a_n) \quad (1.2)$$

for $i_1 < i_2 < \dots < i_k$ and $a_1, \dots, a_{k-1}$ taken from the kernels of the states (if the first unit is in a different configuration of indices, then it vanishes).

This observation leads to a construction of a sequence of states $(\phi^{(m)})_{m=1}^\infty$ on $\sqcup_{i \in I} \mathcal{A}_i$, defined by (1.1) on $\sqcup_{i \in I} \mathcal{A}_i$ and identifying units on an increasing sequence of ‘subspaces’. This gives a discrete interpolation between the monotone product for $m = 1$ and the free product obtained in the limit $m \rightarrow \infty$, which in some sense ‘neglects order’ with increasing m . The sequence of product states is called the *monotone hierarchy* and the associated ‘independent’ variables are called *m-monotone*.

The pace of this interpolation is based on the combinatorics of ordered non-crossing partitions with monotone order (color) starting from depth m , giving monotone non-crossing partitions for $m = 1$ and all ordered non-crossing partitions for $m = \infty$. In particular, the (even) moments of the standard (mean zero, variance one) central limit laws are given by

$$M_{2k}^{(m)} = 1/k! |\mathcal{ONC}_{2k}^2(m)| \quad (1.3)$$

where $\mathcal{ONC}_{2k}^2(m)$ stands for ordered non-crossing pair partitions of a $2k$ -element set, which have a monotone order starting from depth m . In particular, for $m = 1$ we get the moments $M_{2k}^{(1)} = 2^{-k} \binom{2k}{k}$ of the arcsine law supported on $[-\sqrt{2}, \sqrt{2}]$, and in the limit $m \rightarrow \infty$ we obtain the moments $M_{2k}^{(\infty)} = \frac{1}{k+1} \binom{2k}{k}$ of the Wigner measure supported on $[-2, 2]$.

On the level of Cauchy transforms of central limit laws, the interpolation can be nicely described in terms of their continued fractions. Thus, let

$$G(z) = \frac{1}{z - \frac{\beta_1}{z - \frac{\beta_2}{z - \frac{\beta_3}{\ddots}}}} \quad (1.4)$$

be the Cauchy transform of a probability measure associated with the Jacobi sequence $(\beta_n) = (\beta_1, \beta_2, \beta_3, \dots)$. Then, to sequences

$$(\beta_n^{(m)}) = (\underbrace{1, 1, \dots, 1}_{m \text{ times}}, 1/2, 1/2, \dots) \quad (1.5)$$

correspond Cauchy transforms $G^{(m)}(z)$ of the m -monotone central limit measures. In particular, for $m = 1$ we get $(\beta_n^{(1)}) = (1, 1/2, 1/2, \dots)$, corresponding to the arcsine law, and for $m = \infty$, we get $(\beta_n^{(\infty)}) = (1, 1, 1, \dots)$, corresponding to the Wigner law. Thus, successive iterations of $G^{(\infty)}(z)$ are obtained from $G^{(1)}(z)$ by taking the $m - 1$ -th

power of the right-sided shift on the set of Jacobi sequences of the form (1.3), with 1's replacing 1/2's at the first m sites.

We also derive the invariance principle and give a realization of the limits using m -monotone Gaussian operators $\omega^{(m)}(f)$, where $f \in \Theta = \{\chi_{(s,t]} : 0 \leq s < t\}$, on truncations $\mathcal{F}^{(m)}(\mathcal{H})$ of the free Fock space $\mathcal{F}(\mathcal{H})$, where $\mathcal{H} = L^2(\mathbb{R}^+)$, called m -monotone Fock spaces. If $f_1, f_2, \dots, f_{2k} \in \Theta$ have identical or disjoint supports, then explicit combinatorial formulas for the mixed moments of $\omega^{(m)}(f_1), \omega^{(m)}(f_2), \dots, \omega^{(m)}(f_{2k})$ in the vacuum state φ are derived. It is worth noting that even in the monotone case we get a new formula

$$\varphi(\omega^{(1)}(f_1)\omega^{(1)}(f_2)\dots\omega^{(1)}(f_{2k})) = \sum_{\substack{\pi \in \mathcal{NC}_{2k}^2 \\ \pi \sim (f_1, \dots, f_{2k})}} \prod_{i=1}^k \frac{t^{(i)} - s^{(i)}}{\text{Inn}(\pi_i) + 1} \quad (1.6)$$

where summation runs over non-crossing partitions $\pi = \{\pi_1, \pi_2, \dots, \pi_k\}$ ‘compatible’ (in the sense specified in Section 10) with the supports of $f_1, f_2, \dots, f_n$ (support $(s^{(i)}, t^{(i)})$ is associated with block π_i), with $\text{Inn}(\pi_i)$ denoting the number of blocks which are inner with respect to block π_i and are associated with supports identical to that associated with π_i .

Our approach resembles the construction of the *hierarchy of freeness* [L1] (see [FL] for the limit theorems) which gave a discrete interpolation between the boolean product of states and the free product of states. The latter is understood in [L1] in the weak sense of convergence of moments. However, a reduction of free independence (free product of states) to tensor independence (tensor product of states) in the strong sense (reduction of the free product of states to the tensor product of states on extended algebras) has recently also been given [L2]. A similar approach to that in [L2] can also be given for the monotone hierarchy.

Other interpolations between various models of quantum probability were studied in [BLS], [BW].

2. PRODUCTS OF STATES

In this section we introduce a new definition of the monotone product of states, which allows us to compare it with the free product of states and later leads to the construction of a discrete interpolation between the two products.

By a *noncommutative probability space* we understand a pair $(\mathcal{A}, \phi)$, where $\mathcal{A}$ is a unital $*$ -algebra over $\mathbb{C}$ and $\phi : \mathcal{A} \rightarrow \mathbb{C}$ is a positive ($\phi(x^*x) \geq 0$ for all $x \in \mathcal{A}$) normalized ($\phi(1) = 1$) linear functional called *state*. Elements of the $*$ -algebra $\mathcal{A}$ are called *random variables*. A noncommutative probability space $(\mathcal{A}, \phi)$ is called a $\mathcal{C}^*$ -probability space if $\mathcal{A}$ is a unital $\mathcal{C}^*$ -algebra. By the *distribution* of the random variable $a \in \mathcal{A}$ we understand the linear functional $\mu_a : \mathbb{C}[X] \rightarrow \mathbb{C}$, defined on the polynomials of X by

$$\mu_a(P) = \phi(P(a)).$$

Of course, the distribution μ_a is uniquely determined by the moments $\phi(a^n)$ of the variable a for $n \in \mathbb{N}$.

We assume in this paper that I is a linearly ordered set of indices.

DEFINITION 2.1. By the *monotone product* of states ϕ_i we understand the state ϕ on the free product $\sqcup_{i \in I} \mathcal{A}_i$ without identification of units given by

$$\phi(a_1 \dots a_{k-1} a_k a_{k+1} \dots a_n) = \phi_{i_k}(a_k) \phi(a_1 \dots a_{k-1} a_{k+1} \dots a_n), \quad (2.1)$$

for $a_j \in \mathcal{A}_{i_j}$ whenever neighboring variables come from different algebras and $i_{k-1} < i_k > i_{k+1}$ (with only one inequality if $k \in \{1, n\}$).

Note that the freeness condition (1.1) is of quite different form than the monotonicity condition given by (2.1). However, as we demonstrate below, one can reduce Definition 2.1 to the freeness condition plus conditions on moments involving units. In the case of the free product of states, the condition on moments involving units is trivial, namely units of $*$ -algebras $\mathcal{A}_i$ always play the role of a ‘global unit’ and can be identified. In the case of the monotone product of states, units of $*$ -algebras $\mathcal{A}_i$ play the role of a ‘monotone approximate unit’. This can be done even if $*$ -algebras $\mathcal{A}_i$ are not unital - it is enough to take their unitizations.

DEFINITION 2.2. Let $(\mathcal{A}_i, \phi_i)_{i \in I}$ be a family of noncommutative probability spaces. Define $\psi : \sqcup_{i \in I} \mathcal{A}_i \rightarrow \mathbb{C}$ to be the linear functional defined by the freeness condition (1.1) for all a_j in the kernels of the ϕ_{i_j} , respectively, and by the condition

$$\psi(a_1 \dots a_{k-1} 1_{i_k} a_{k+1} \dots a_n) = \begin{cases} \psi(a_1 \dots a_{k-1} a_{k+1} \dots a_n) & \text{if } i_1 < i_2 < \dots < i_k \\ 0 & \text{otherwise} \end{cases} \quad (2.2)$$

where $1 \leq k \leq n$ and only the variables a_j preceding 1_{i_k} are assumed to be in the kernels of the states ϕ_{i_j} , respectively, with $\phi_i = \psi|_{\mathcal{A}_i}$ for every $i \in I$.

Example 2.1. Let $I = \{1, 2\}$ (with natural order) and $a \in \mathcal{A}_1$, $b \in \mathcal{A}_2$ and denote $a^0 = a - \psi(a)1_1$, $b^0 = b - \psi(b)1_2$. Then $\psi(a^0 1_2 a^0) = \psi((a^0)^2)$ and $\psi(b^0 1_1 b^0) = 0$ are the key formulas which, together with (1.1), lead to different factorizations

$$\begin{aligned} \psi(aba) &= \psi(a^2)\psi(b) \\ \psi(bab) &= \psi^2(b)\psi(a) \end{aligned}$$

characteristic of the monotone product of states.

THEOREM 2.3. *The state ψ agrees with the monotone product of states ϕ_i , $i \in I$.*

Proof. Let $a_1 \in \mathcal{A}_{i_1}$, $a_2 \in \mathcal{A}_{i_2}, \dots, a_n \in \mathcal{A}_{i_n}$. Denote $a_j^0 = a_j - \phi_{i_j}(a_j)1_{i_j}$ for any $a_j \in \mathcal{A}_{i_j}$ and let ϕ stand for the monotone product of states ϕ_i . Using (2.2) for $n = 1$ and the fact that $\psi(1_{i_1}) = 1$, we obtain

$$\psi(a_1) = \psi(a_1^0 + \phi_{i_1}(a_1)1_{i_1}) = \phi_{i_1}(a_1).$$

Suppose now that ψ agrees with the monotone product of ϕ_i 's on words of length smaller than n and assume that $i_1 < i_2 < \dots < i_k > i_{k+1}$ (if $k = n$, then the last inequality is ignored). Decomposing $a_j = a_j^0 + \phi(a_j)1_{i_j}$ for $1 \leq j \leq k$, we obtain

$$\begin{aligned}\psi(a_1 a_2 \dots a_n) &= \psi(a_1^0 \dots a_k^0 a_{k+1} \dots a_n) \\ &+ \sum_{1 \leq j \leq k} \psi(a_1^0 \dots 1_{i_j} a_{j+1}^0 \dots a_k^0 a_{k+1} \dots a_n) \\ &+ \sum_{1 \leq j < r \leq k} \psi(a_1^0 \dots 1_{i_j} a_{j+1}^0 \dots 1_{i_r} a_{r+1}^0 \dots a_k^0 a_{k+1} \dots a_n) \\ &+ \dots \\ &+ \psi(1_{i_1} \dots 1_{i_k} a_{k+1} \dots a_n).\end{aligned}$$

Now, applying (2.2) to all moments except the first one, we can reduce them to moments of order $< n$. Therefore, by the inductive assumption, they agree with the corresponding moments in the state ϕ . A typical calculation is of the form

$$\begin{aligned}\psi(a_1^0 \dots 1_{i_j} a_{j+1}^0 \dots a_n) &= \psi(a_1^0 \dots a_{j+1}^0 \dots a_n) \\ &= \phi(a_1^0 \dots a_{j+1}^0 \dots a_n) \\ &= \phi(a_1^0 \dots 1_{i_j} a_{j+1}^0 \dots a_n)\end{aligned}$$

where the last equation is an easy property of ϕ for $i_1 < i_2 < \dots < i_k$. Thus we get

$$\begin{aligned}\psi(a_1 a_2 \dots a_n) &= \psi(a_1^0 a_2^0 \dots a_k^0 a_{k+1} \dots a_n) \\ &+ \phi(a_1 a_2 \dots a_n) - \phi(a_1^0 a_2^0 \dots a_k^0 a_{k+1} \dots a_n).\end{aligned}$$

However, from (2.1) we obtain

$$\phi(a_1^0 a_2^0 \dots a_k^0 a_{k+1} \dots a_n) = 0$$

since $a_k^0 \in \text{Ker} \phi_{i_k}$ and $i_{k-1} < i_k > i_{k+1}$. In turn, using (2.2) again we get

$$\begin{aligned}\psi(a_1^0 a_2^0 \dots a_k^0 a_{k+1} \dots a_n) &= \psi(a_1^0 a_2^0 \dots a_k^0 (a_{k+1}^0 + 1_{i_{k+1}}) \dots a_n) \\ &= \psi(a_1^0 a_2^0 \dots a_k^0 a_{k+1}^0 a_{k+2} \dots a_n) \\ &= \dots \\ &= \psi(a_1^0 a_2^0 \dots a_n^0) \\ &= 0\end{aligned}$$

which completes the proof of the inductive step. This allows us to pull out the moment corresponding to the first local maximum in the sequence $(i_1, \dots, i_n)$. In the same manner we can pull out the moment corresponding to every local maximum. ■

3. THE MONOTONE HIERARCHY

In this section we introduce a sequence of product states indexed by $m \in \mathbb{N}$, which gives a discrete interpolation between the monotone product of states ($m = 1$) and the free product of states ($m = \infty$).

DEFINITION 3.1. Let $\{\mathcal{A}_i\}_{i \in I}$ be a family of unital $*$ -algebras with units $\{1_i\}_{i \in I}$, where I is a linearly ordered set, and let $\phi : \sqcup_{i \in I} \mathcal{A}_i \rightarrow \mathbb{C}$ be a linear functional. The family $\{1_i\}_{i \in I}$ is called an *m-monotone family of units* with respect to ϕ if $\phi(1_i) = 1$ for all $i \in I$ and

$$\phi(a_1 \dots a_{j-1} 1_{i_j} a_{j+1} \dots a_n) = \phi(a_1 \dots a_{j-1} a_{j+1} \dots a_n)$$

whenever $j \leq m$ or ($m < j \leq r$ and $i_1 \neq \dots \neq i_m < \dots < i_r > i_{r+1}$) and otherwise the moment vanishes, where $a_1 \in \mathcal{A}_{i_1} \cap \text{Ker } \phi, \dots, a_{j-1} \in \mathcal{A}_{i_{j-1}} \cap \text{Ker } \phi$ and $a_{j+1} \in \mathcal{A}_{i_{j+1}}, \dots, a_n \in \mathcal{A}_{i_n}$.

Let us observe that if $\{1_i\}_{i \in I}$ is an *m-monotone family of units* w.r.t. ϕ , then each 1_i behaves as a unit when it appears at the first m positions in a given moment or if its position precedes the first disorder in the sequence $(i_m, \dots, i_n)$. Otherwise, 1_i acts as a null projection.

DEFINITION 3.2. Let $m \in \mathbb{N}$ and let $\{(\mathcal{A}_i, \phi_i)\}_{i \in I}$ be a family of noncommutative probability spaces, where I is a linearly ordered index set. By the *m-monotone product of states* $\phi_i, i \in I$, we understand the linear functional $\phi : \sqcup_{i \in I} \mathcal{A}_i \rightarrow \mathbb{C}$ such that

$$\phi(a_1 a_2 \dots a_n) = 0, \quad \text{whenever } a_k \in \mathcal{A}_{i_k} \cap \text{Ker } \phi_{i_k}$$

and $\{1_i\}_{i \in I}$ is an *m-monotone family of units* w.r.t. ϕ . The sequence of *m-monotone product states* will be called the *monotone hierarchy (of product states)*.

With the *m-monotone product of states* we can associate a notion of ‘independence’ in a natural way for every m .

DEFINITION 3.3. Let $\{\mathcal{A}_i\}_{i \in I}$ be a family of $*$ -subalgebras of a noncommutative probability space $(\mathcal{A}, \phi)$. A family of projections $\{1_i \in \mathcal{A}_i\}_{i \in I}$ is an *m-monotone family of units w.r.t. ϕ* if 1_i is an internal unit in $\mathcal{A}_i$ for every $i \in I$ and conditions of Definition 3.1 are satisfied. If such a family exists, then we say that $\{\mathcal{A}_i\}$ are *m-monotone* with respect to ϕ if and only if

$$\phi(a_1 a_2 \dots a_n) = 0 \quad \text{whenever } a_k \in \mathcal{A}_{i_k} \cap \text{Ker } \phi$$

where $k = 1, \dots, n$ and $i_1 \neq \dots \neq i_n$. The associated sequence of independences will be called the *monotone hierarchy (of freeness)*.

In particular, $*$ -subalgebras are 1-monotone w.r.t. ϕ if and only if they are monotone independent w.r.t. ϕ . Also, if we extend Definitions 3.1 and 3.3 to include the case $m = \infty$, we can say that $*$ -subalgebras are ∞ -monotone w.r.t. ϕ if and only if they are

free w.r.t. ϕ . In turn, the m -monotone hierarchy contains models which give examples of ‘freeness without identification of units’.

Example 3.1. Let $\mathcal{H}$ be a Hilbert space, $\Omega \in \mathcal{H}$ - a unit vector and $\varphi(\cdot) = \langle \cdot, \Omega \rangle$ - the associated state. Then $1_1 = 1 \otimes P_\Omega$, $1_2 = 1 \otimes 1$ give a 1-monotone family of units w.r.t. the state $\varphi \otimes \varphi$, with respect to which $\mathcal{A}_1 = B(\mathcal{H}) \otimes P_\Omega$ and $\mathcal{A}_2 = 1 \otimes B(\mathcal{H})$ are 1-monotone.

LEMMA 3.4. *Let $\{\mathcal{A}_i\}_{i \in I}$ be a family of $*$ -subalgebras of a noncommutative probability space $(\mathcal{A}, \phi)$ which are m -monotone with respect to ϕ . Then*

$$\phi(a_1^0 \dots a_r^0 a_{r+1} \dots a_n) = 0$$

whenever $i_1 \neq \dots \neq i_m < i_{m+1} < \dots < i_r > i_{r+1}$, where $a_k^0 \in \mathcal{A}_{i_k} \cap \text{Ker} \phi$ for $1 \leq k \leq m$ and $a_r \in \mathcal{A}_{i_r}$ for $m+1 \leq r \leq n$.

Proof. Using the definition of the m -monotone family of units, we obtain

$$\begin{aligned} \phi(a_1^0 \dots a_r^0 a_{r+1} \dots a_n) &= \phi(a_1^0 \dots a_r^0 a_{r+1}^0 \dots a_n) + \phi(a_{r+1}) \phi(a_1^0 \dots a_r^0 1_{\mathcal{A}_{i_{r+1}}} a_{r+2} \dots a_n) \\ &= \phi(a_1^0 \dots a_r^0 a_{r+1}^0 a_{r+2} \dots a_n) \end{aligned}$$

We repeat the same for $a_{r+2}, \dots, a_n$, which leads to

$$\phi(a_1^0 \dots a_r^0 a_{r+1} \dots a_n) = \phi(a_1^0 \dots a_r^0 a_n^0)$$

and this vanishes by the definition of m -monotone $*$ -subalgebras. ■

LEMMA 3.5. *Let $\{\mathcal{A}_i\}_{i \in I}$ be a family of $*$ -subalgebras of a noncommutative probability space $(\mathcal{A}, \phi)$ which are m -monotone with respect to ϕ . Then*

$$\phi(a_1 \dots a_{j-1} 1_{i_j} a_{j+1} \dots a_n) = \phi(a_1 \dots a_{j-1} a_{j+1} \dots a_n)$$

whenever $j = 1, \dots, r$ and $i_1 \neq \dots \neq i_m < i_{m+1} < \dots < i_r > i_{r+1}$, where $a_k \in \mathcal{A}_{i_k}$ for $1 \leq k \leq n$.

Proof. For $n = 1$ the statement is obvious. Suppose it holds for $1 \leq s \leq n - 1$. Then, decomposing $a_k = a_k^0 + \phi(a_k) 1_{i_k}$ succesively for $k = 1, \dots, j - 1$, we obtain

$$\begin{aligned} \phi(a_1 \dots a_{j-1} 1_{i_j} a_{j+1} \dots a_n) &= \sum_{k=1}^{j-1} \phi(a_k) \phi(a_1^0 \dots a_{k-1}^0 a_{k+1} \dots a_{j-1} 1_{i_j} a_{j+1} \dots a_n) \\ &\quad + \phi(a_1^0 \dots a_{j-1}^0 1_{i_j} a_{j+1} \dots a_n) \end{aligned}$$

which, by the inductive assumption and the definition of the m -monotone family of units, gives

$$\sum_{k=1}^{j-1} \phi(a_k) \phi(a_1^0 \dots a_{k-1}^0 a_{k+1} \dots a_{j-1} a_{j+1} \dots a_n) + \phi(a_1^0 \dots a_{j-1}^0 a_{j+1} \dots a_n)$$

$$= \phi(a_1 \dots a_{j-1} a_{j+1} \dots a_n)$$

and that completes the induction proof. $\blacksquare$.

Example 3.2. Suppose that $\mathcal{A}_1, \mathcal{A}_2$ are 2-monotone $*$ -subalgebras of $(\mathcal{A}, \phi)$ and $\{1_1, 1_2\}$ is a 2-monotone family of units. Let $a \in \mathcal{A}_1, b \in \mathcal{A}_2$. It can be easily shown that mixed moments of orders ≤ 4 agree with the corresponding moments in the free case. Namely

$$\begin{aligned} \phi(ab) &= \phi(a)\phi(b), \\ \phi(aba) &= \phi(a^2)\phi(b), \\ \phi(abab) &= \phi(a^2)\phi^2(b) + \phi(b^2)\phi^2(a) - \phi^2(a)\phi^2(b). \end{aligned}$$

Let us compute the ‘alternating’ mixed moment of order 5. We have

$$\begin{aligned} \phi(ababa) &= \phi(a^0 b a b a) + \phi(a)\phi(b a b a) \\ &= \phi(a^0 b^0 a b a) + \phi(b)\phi(a^0 a b a) + \phi(a)\phi(b^0 a b a) + \phi(a)\phi(b)\phi(a b a). \end{aligned}$$

Let us observe that the first term in this expression vanishes by Lemma 3.4. Besides, easy computations of lower-order moments give

$$\begin{aligned} \phi(a^0 a b a) &= \phi(b)\phi(a^3) - \phi(b)\phi(a)\phi(a^2), \\ \phi(b^0 a b a) &= \phi^2(a)\phi(b^2) - \phi^2(a)\phi^2(b), \\ \phi(a b a) &= \phi(b)\phi(a^2). \end{aligned}$$

Thus we finally obtain

$$\phi(ababa) = \phi^3(a)\phi(b^2) - \phi^3(a)\phi^2(b) + \phi(a^3)\phi^2(b),$$

which gives the lowest order moment for $m = 2$ which differs from the corresponding moments in both the free and the monotone cases.

4. HILBERT SPACE REPRESENTATIONS

We begin this section with recalling the definition of the free product of noncommutative probability spaces [V]. Let I be a linearly ordered set of indices and let $(\mathcal{A}_i, \phi_i)_{i \in I}$ be a family of $\mathcal{C}^*$ -noncommutative probability spaces. For each $i \in I$ we denote by $(\mathcal{H}_i, \pi_i, \xi_i)$ the GNS triple of $(\mathcal{A}_i, \phi_i)$, i.e. $\mathcal{H}_i$ is a Hilbert space, ξ_i is a cyclic (unit) vector in $\mathcal{H}_i$ and $\pi_i : \mathcal{A}_i \rightarrow \mathcal{B}(\mathcal{H}_i)$ is a $*$ -homomorphism, such that

$$\phi_i(a) = \langle \pi_i(a)\xi_i, \xi_i \rangle$$

for all $a \in \mathcal{A}_i$, where $\langle \cdot, \cdot \rangle$ denotes the scalar product in $\mathcal{H}_i$ (for simplicity, we use the same notation for all scalar products).

Let $\mathcal{H}_i^0$ be the orthogonal complement in $\mathcal{H}_i$ of the one-dimensional subspace generated by ξ_i , i.e. $\mathcal{H}_i^0 = \mathcal{H}_i \ominus \mathbb{C}\omega_i$. For any $h \in \mathcal{H}_i$, let h^0 denote the orthogonal projection of h on $\mathcal{H}_i^0$. Besides, let $\mathcal{H}$ denote the Hilbert space

$$\mathcal{H} = \mathbb{C}\Omega \oplus \bigoplus_{n=1}^{\infty} \bigoplus_{i_1 \neq i_2 \neq \dots \neq i_n} \mathcal{H}_{i_1}^0 \otimes \mathcal{H}_{i_2}^0 \otimes \dots \otimes \mathcal{H}_{i_n}^0 \quad (4.1)$$

with the canonical scalar product, where Ω is the so-called vacuum (unit) vector. The pair $(\mathcal{H}, \Omega)$ is called the *free product* of $(\mathcal{H}_i, \xi_i)$.

On $\mathcal{H}$ we define a $*$ -representation $\lambda_i : \mathcal{A}_i \rightarrow \mathcal{B}(\mathcal{H})$ of each algebra $\mathcal{A}_i$, $i \in I$ as follows:

$$\begin{aligned} \lambda_i(a)\Omega &= (\pi_i(a)\xi_i)^0 + \langle \pi_i(a)\xi_i, \xi_i \rangle \Omega \\ \lambda_i(a)(h_1 \otimes \dots \otimes h_n) &= \begin{cases} (\pi_i(a)\xi_i)^0 \otimes h_1 \otimes \dots \otimes h_n + \langle \pi_i(a)\xi_i, \xi_i \rangle h_1 \otimes \dots \otimes h_n & \text{if } i \neq i_1 \\ (\pi_i(a)h_1)^0 \otimes h_2 \otimes \dots \otimes h_n + \langle \pi_i(a)h_1, \xi_i \rangle h_2 \otimes \dots \otimes h_n & \text{if } i = i_1 \end{cases} \end{aligned}$$

Finally, on $\mathcal{B}(\mathcal{H})$ we define the so-called *vacuum state* $\varphi(\cdot) = \langle \cdot, \Omega \rangle$.

In order to introduce the m -monotone product of Hilbert spaces, we want to characterize admissible simple tensors which appear in the product. For that purpose we introduce the associated families of sequences $I_n(m) \subset I^n$. We set

$$I_n(m) = \{(i_1, \dots, i_n); i_1 \neq i_2 \neq \dots \neq i_n\} \text{ for } n \leq m,$$

whereas

$$I_n(m) = \{(i_1, \dots, i_n); i_1 > \dots > i_{n-m+1} \neq i_{n-m+2} \neq \dots \neq i_n\} \text{ for } n > m.$$

Thus, in contrast to the free product case, all sequences $(i_1, i_2, \dots, i_n)$ subject to $i_1 \neq i_2 \neq \dots \neq i_n$ are allowed only if $n \leq m$. In turn, if $n > m$, then the first $n - m + 1$ indices must be in a decreasing order.

DEFINITION 4.1. Let $\mathcal{H}^{(m)}$ be the closed subspace of the Hilbert space $\mathcal{H}$ of the form

$$\mathcal{H}^{(m)} = \mathbb{C}\Omega \oplus \bigoplus_{n=1}^{\infty} \bigoplus_{(i_1, \dots, i_n) \in I_n(m)} \mathcal{H}_{i_1}^0 \otimes \mathcal{H}_{i_2}^0 \otimes \dots \otimes \mathcal{H}_{i_n}^0 \quad (4.2)$$

The pair $(\mathcal{H}^{(m)}, \Omega)$ will be called the *m -monotone product of Hilbert spaces* $(\mathcal{H}_i, \xi_i)$.

It is natural to compare this truncation with the *m -free product of Hilbert spaces* [F-L] given by

$$\mathcal{H}_{(m)} = \mathbb{C}\Omega \oplus \bigoplus_{n=1}^m \bigoplus_{i_1 \neq i_2 \neq \dots \neq i_n} \mathcal{H}_{i_1}^0 \otimes \mathcal{H}_{i_2}^0 \otimes \dots \otimes \mathcal{H}_{i_n}^0$$

and observe that, in contrast to $\mathcal{H}^{(m)}$, $\mathcal{H}_{(m)}$ is a truncation of the free product of Hilbert spaces which takes into account only sequences $(i_1, i_2, \dots, i_n)$ of length $\leq m$. In the case of $\mathcal{H}^{(m)}$, the truncation is more delicate and depends on the order in which indices appear. Thus, for instance, if I is infinite, simple tensors of arbitrary length appear in $\mathcal{H}^{(m)}$.

Example 4.1. Let $I = \{1, 2\}$. If $m = 1$, we obtain

$$\mathcal{H}^{(1)} = \mathbb{C}\Omega \oplus \mathcal{H}_1^0 \oplus \mathcal{H}_2^0 \oplus (\mathcal{H}_2^0 \otimes \mathcal{H}_1^0)$$

thus the 1-monotone product of $\mathcal{H}_1$ and $\mathcal{H}_2$ coincides with the monotone Fock space. In turn, the 2-monotone product of $\mathcal{H}_1$ and $\mathcal{H}_2$ is of the form

$$\mathcal{H}^{(2)} = \mathbb{C}\Omega \oplus \mathcal{H}_1^0 \oplus \mathcal{H}_2^0 \oplus (\mathcal{H}_2^0 \otimes \mathcal{H}_1^0) \oplus (\mathcal{H}_1^0 \otimes \mathcal{H}_2^0) \oplus (\mathcal{H}_2^0 \otimes \mathcal{H}_1^0 \otimes \mathcal{H}_2^0)$$

thus it is a direct sum of all simple tensors of order ≤ 2 and only one simple tensor of order 3.

Let us introduce the closed subspace of $\mathcal{H}^{(m)}$ given by

$$\mathcal{H}^{(m)}(i) = \mathcal{H}_{(m-1)} \oplus \left(\bigoplus_{n=m}^{\infty} \bigoplus_{\substack{(i_1, \dots, i_n) \in I_n(m) \\ i_1 \leq i}} \mathcal{H}_{i_1}^0 \otimes \dots \otimes \mathcal{H}_{i_n}^0 \right),$$

where $i \in I$, which allows us to define suitable truncations of the free product representation. Let representations $\lambda_i^{(m)}$ of $\mathcal{A}_i$ on $\mathcal{H}^{(m)}$ be defined by

$$\lambda_i^{(m)} : \mathcal{A}_i \rightarrow \mathcal{B}(\mathcal{H}^{(m)}) \quad , \quad \lambda_i^{(m)}(a)h = \begin{cases} \lambda_i(a)h & \text{if } h \in \mathcal{H}^{(m)}(i) \\ 0 & \text{if } h \in \mathcal{H}^{(m)}(i)^\perp \end{cases}.$$

Thus, $\lambda_i^{(m)}$ is a truncation of λ_i for every m . The ‘level’ of this truncation depends on the order of Hilbert spaces at all sites except the last m sites where the order is irrelevant as in the case of representations λ_i . Of course, the free product $*_{i \in I} \lambda_i^{(m)}$ is then a truncation of the free product representation $*_{i \in I} \lambda_i$ and agrees with the latter on the m -free product of Hilbert spaces.

THEOREM 4.2. *Algebras $\{\lambda_i^{(m)}(\mathcal{A}_i)\}_{i \in I}$ are m -monotone $*$ -subalgebras of the non-commutative probability space $(\mathcal{B}(\mathcal{H}^{(m)}), \varphi)$, where φ is the vacuum state on $\mathcal{B}(\mathcal{H}^{(m)})$.*

Proof. Let us fix $m \in \mathbb{N}$. We will show that $\{\lambda_i^{(m)}(1_i)\}_{i \in I}$ is an m -monotone family of units for $\{\lambda_i^{(m)}(\mathcal{A}_i)\}_{i \in I}$. Notice that

$$\lambda_i^{(m)}(1_i)h = \begin{cases} h & \text{if } h \in \mathcal{H}^{(m)}(i) \\ 0 & \text{if } h \in \mathcal{H}^{(m)}(i)^\perp \end{cases}.$$

Besides, let $a_1 \in \mathcal{A}_{i_1}, \dots, a_n \in \mathcal{A}_{i_n}$ be such that $i_1 \neq i_2 \neq \dots \neq i_n$. For notational simplicity we denote $X_k = \lambda_{i_k}^{(m)}(a_{i_k})$. Then

$$\begin{aligned} \varphi(X_1 \dots X_{j-1} \lambda_i^{(m)}(1_{i_j}) X_{j+1} \dots X_n) &= \langle X_1 \dots X_{j-1} \lambda_i^{(m)}(1_{i_j}) X_{j+1} \dots X_n \Omega, \Omega \rangle \\ &= \langle \Omega, X_n^* \dots X_{j+1}^* \lambda_i^{(m)}(1_{i_j}) X_{j-1}^* \dots X_1^* \Omega \rangle. \end{aligned}$$

Let $X_k \in \text{Ker} \phi$ for $k = 1, \dots, j-1$. If $(i_j, \dots, i_1) \in I_j(m)$, then it can be seen that $X_{j-1}^* \dots X_1^* \Omega$ is a simple tensor from $\mathcal{H}^{(m)}(i_j)$, hence

$$\begin{aligned} \varphi(X_1 \dots X_{j-1} \lambda_i^{(m)}(1_{i_j}) X_{j+1} \dots X_n) &= \langle \Omega, X_n^* \dots X_{j+1}^* \lambda_i^{(m)}(1_{i_j}) X_{j-1}^* \dots X_1^* \Omega \rangle \\ &= \langle \Omega, X_n^* \dots X_{j+1}^* X_{j-1}^* \dots X_1^* \Omega \rangle \\ &= \varphi(X_1 \dots X_{j-1} X_{j+1} \dots X_n). \end{aligned}$$

If, in turn, $(i_j, \dots, i_1) \notin I_j(m)$, then $X_{j-1}^* \dots X_1^* \Omega \in \mathcal{H}^{(m)}(i_j)^\perp$, hence

$$\lambda_i^{(m)}(1_{i_j}) X_{j-1}^* \dots X_1^* \Omega = 0,$$

which proves that $\{\lambda_i^{(m)}(1_i)\}_{i \in I}$ is an m -monotone family of units.

Let us show now that variables $X_1, \dots, X_n$ satisfy conditions of Definition 3.3. Assume that $X_1, \dots, X_n \in \text{Ker} \phi$. Then $X_n^* \dots X_1^* \Omega \perp \mathbb{C} \Omega$ since it is a simple tensor from $\mathcal{H}^{(m)}(i_n)$, and thus

$$\varphi(X_1 \dots X_n) = \langle \Omega, X_n^* \dots X_1^* \Omega \rangle = 0,$$

which completes the proof. ■

COROLLARY 4.3. *The m -monotone product of states is a state which agrees with the free product of states on words of length less than or equal to $2m$.*

Proof. It follows from the GNS representation in the proof of Theorem 4.2 that the m -monotone product of states is positive, thus it is a state. Moreover, by definition, it satisfies the freeness condition and the units act as an identified unit at the first m places in the moment. From this and the fact that any state is a hermitian functional it follows that units act as an identified unit at the last m places in the moment. Therefore, the moments have to agree if their lengths are smaller or equal to $2m$. ■

5. COMBINATORICS

This section is devoted to the combinatorics of the monotone hierarchy. It is based on ordered non-crossing partitions with disorders starting at different depths (compare with the combinatorics of non-crossing partitions in free probability [S]).

An *ordered partition* P of the set $\{1, 2, \dots, n\}$ is a tuple $(P_1, P_2, \dots, P_k)$ of disjoint non-empty subsets called *blocks* such that $P_1 \cup P_2 \cup \dots \cup P_k = \{1, 2, \dots, n\}$. For any block P_r the number r is the *order* of P_r , which can also be interpreted as *color*. If blocks are two-element sets, then P is called an *ordered pair-partition*. An ordered

partition S is called *crossing* if and only if there exist $s, s' \in P_i$ and $r, r' \in P_j$ for some $i \neq j$ such that $s < r < s' < r'$. If P is not a crossing partition, then it is called *non-crossing*. The collection of ordered non-crossing partitions (pair-partitions) of the set $\{1, 2, \dots, n\}$ will be denoted $\mathcal{ONC}_n$ ($\mathcal{ONC}_n^2$). Block P_i is called *inner* with respect to block P_j , which we denote $P_j < P_i$, if and only if there exist two numbers $p, q \in P_j$ such that the interval $(p, q) \cap P_i = \emptyset$ and for every $s \in P_i$ we have $p < s < q$. Then P_j will be called *outer* with respect to P_i . By the *depth* of block P_i , denoted $d(P_i)$, we will understand the number of blocks of P which are outer w.r.t. P_i or coincide with P_i . In particular, if there are no blocks which are outer w.r.t. P_i , then $d(P_i) = 1$.

DEFINITION 5.1. By $\mathcal{ONC}_n(m)$ we will denote the collection of all *ordered non-crossing partitions* $P = (P_1, P_2, \dots, P_k)$ of the set $\{1, 2, \dots, n\}$ such that for all $i, j \in \{1, \dots, k\}$ the following implication holds:

$$d(P_j) \geq m \text{ and } P_j < P_i \implies j < i.$$

In other words, the color of blocks is a monotone function of their depths starting from depth m . In particular, we call $\mathcal{ONC}_n(1)$ *monotone non-crossing partitions* (we also denote them $\mathcal{MNC}_n$). The corresponding pair partitions will be denoted $\mathcal{ONC}_n^2(m)$ and $\mathcal{MNC}_n^2$, respectively.

Example 5.1. For instance, for $n = 8$ the tuples

$$P = (\{4, 7\}, \{1, 8\}, \{2, 3\}, \{5, 6\}) \text{ and } R = (\{1, 8\}, \{2, 3\}, \{4, 7\}, \{5, 6\})$$

are different ordered pair-partitions of the set $\{1, 2, \dots, 8\}$. The corresponding diagrams are drawn in Figure 1, where blocks are labelled by colors associated with their orders.




Figure 1. Diagrams of partitions P and R .

In the case of partitions P and R in Figure 1 all blocks but $\{1, 8\}$ are inner w.r.t. $\{1, 8\}$, but $\{5, 6\}$ is the only block which is inner w.r.t. $\{4, 7\}$. The depths are given by $d(\{1, 8\}) = 1$, $d(\{2, 3\}) = d(\{4, 7\}) = 2$, $d(\{5, 6\}) = 3$. Note that $R \in \mathcal{MNC}_8^2$ since its coloring is monotone, whereas $P \in \mathcal{ONC}_8^2(2) \setminus \mathcal{MNC}_8^2$ since the coloring of the whole partition is not monotone, however it is monotone starting from depth 2.

PROPOSITION 5.2. For every $k \in \mathbb{N}$ we have $|\mathcal{MNC}_{2k}^2| = (2k - 1)!!$.

Proof. The proof is elementary and is left to the reader. ■

DEFINITION 5.3. Let $i_1, \dots, i_n \in \mathbb{N}$ and set $\{i_1, \dots, i_n\} = \{k_1, \dots, k_r\}$, where $k_1 < k_2 < \dots < k_r$. By an ordered partition *associated* with the tuple $(i_1, \dots, i_n)$ we understand the partition $P = (P_1, \dots, P_r)$ given by $P_j = \{s : i_s = k_j\}$. We then write $P \sim (i_1, \dots, i_n)$.

Example 5.2. Let $i_1 = i_4 = 2$, $i_2 = i_5 = 4$, $i_3 = 1$. Then $(P_1, P_2, P_3) \sim (2, 4, 1, 2, 4)$, where $P_1 = \{3\}$, $P_2 = \{1, 4\}$, $P_3 = \{2, 5\}$.

We shall also need a 'continuous' version of Definition 5.3 with the tuple of indices replaced by a tuple of characteristic functions. For that purpose, on the set

$$\Theta = \{\chi_{(s,t]} : 0 \leq s < t < \infty\} \quad (5.1)$$

introduce a partial order by setting $f = \chi_{(s_1,t_1]} < \chi_{(s_2,t_2]} = g$ if and only if $t_1 \leq s_2$ and $f \leq g$ iff $f < g$ or $f = g$.

DEFINITION 5.4. Let $f_1, f_2, \dots, f_n \in \Theta$ have identical or disjoint supports. We say that $P \in \mathcal{OP}_n$ is *compatible* with the tuple $(f_1, f_1, \dots, f_n)$ if and only if

1. $i, j \in P_k \implies f_i = f_j$,
2. $i \in P_k, j \in P_l$ and $k < l \implies f_i \leq f_j$.

We then write $P \sim (f_1, f_2, \dots, f_n)$.

Example 5.3. Let $f_1 = f_2 = f_5 = f_6 = \chi_{(0,1]}$ and $f_3 = f_4 = \chi_{(1,2]}$. Then the partition $P = (\{1, 6\}, \{2, 5\}, \{3, 4\})$ is compatible with the tuple $(f_1, f_2, \dots, f_6)$, whereas $P' = (\{3, 4\}, \{1, 6\}, \{2, 5\})$ is not since, for instance, block $\{3, 4\}$ precedes block $\{1, 6\}$ and $f_3 = f_4 > f_1 = f_6$.

The combinatorial formulas derived in this paper can be expressed in terms of the usual non-crossing partitions. If $P = (P_1, P_2, \dots, P_k) \in \mathcal{ONC}_n$ and $\pi = \{\pi_1, \dots, \pi_k\} \in \mathcal{NC}_n$ have the same blocks, then we shall write $\pi \sim P$ (by abuse of notation).

Let us assume now that we have a pair of objects: a partition $\pi = \{\pi_1, \pi_2, \dots, \pi_k\} \in \mathcal{NC}_{2k}^2$ and a tuple of functions $(f_1, f_2, \dots, f_{2k})$, where $f_i = \chi_{(s_i, t_i]}$, $i = 1, \dots, 2k$, have identical or disjoint supports and such that $\pi \sim P \sim (f_1, f_2, \dots, f_{2k})$ for some ordered partition P . This implies that if $\{p, q\} = \pi_i$ for some i , then $f_p = f_q := f^{(i)} = \chi_{(s^{(i)}, t^{(i)})}$. Thus $f^{(1)}, f^{(2)}, \dots, f^{(k)}$ will denote characteristic functions associated with blocks $\pi_1, \pi_2, \dots, \pi_k$ and $s^{(1)}, s^{(2)}, \dots, s^{(k)}$ as well as $t^{(1)}, t^{(2)}, \dots, t^{(k)}$ will be the left and the right endpoints of their supports, respectively.

Computations will be based on counting inner blocks which are 'compatible' with the supports of associated characteristic functions. For that purpose, to every block π_i of π we assign the number of its blocks which are inner w.r.t. π_i and are associated with functions of the same supports as $f^{(i)}$, namely

$$\text{Inn}(\pi_i) = \#(\pi_j : \pi_j > \pi_i \text{ and } f^{(j)} = f^{(i)}). \quad (5.2)$$

Note that $\text{Inn}(\pi_i)$ depends on the tuple $(f_1, f_2, \dots, f_{2k})$, but this fact is suppressed in the notation.

Finally, to every π and $(f_1, f_2, \dots, f_{2k})$ we associate the number

$$c_\pi(f_1, f_2, \dots, f_{2k}) = \#(P \in \mathcal{MNC}_{2k}^2 : \pi \sim P \sim (f_1, f_2, \dots, f_{2k})) \quad (5.3)$$

which gives the number of ‘admissible’ colorings of π .

Example 5.4. Let $\pi \in \mathcal{NC}_8^2$ be given by the diagram in Figure 2 with supports $f = \chi_{(0,t]}$, and $g = \chi_{(t,t']}]$ associated with blocks as shown. Set $\pi_1 = \{1, 8\}$, $\pi_2 = \{2, 3\}$, $\pi_3 = \{4, 7\}$, $\pi_4 = \{5, 6\}$.




Figure 2. Diagram of the partition π .

Block π_1 has one inner block with support equal to f , thus $\text{Inn}(\pi_1) = 1$, whereas the remaining blocks have no inner blocks with identical supports, thus $\text{Inn}(\pi_2) = \text{Inn}(\pi_3) = \text{Inn}(\pi_4) = 0$. Note also that $c_\pi(f, g, g, f, g, g, f, f) = 3$ (block π_1 must be colored by 1, block π_2 can be colored by 2, 3 or 4 and for every such choice there is only one coloring of the remaining blocks since $\pi_3 < \pi_4$).

6. LEMMAS ON MIXED MOMENTS

In the lemmas given below we assume that $\{\mathcal{A}_i\}_{i \in I}$ is a family of $*$ -subalgebras of a noncommutative probability space $(\mathcal{A}, \phi)$ which are m -monotone with respect to ϕ and that $a_k \in \mathcal{A}_{i_k}$ for $1 \leq k \leq n$.

LEMMA 6.1. *Let $\{\mathcal{A}_i\}_{i \in I}$ be m -monotone $*$ -subalgebras of $(\mathcal{A}, \phi)$ and let $a_1 \in \mathcal{A}_{i_1}, \dots, a_n \in \mathcal{A}_{i_n}$ be such that $i_1 \neq i_2 \neq \dots \neq i_n$. If there exists $j \in \{1, \dots, n\}$, for which $\phi(a_j) = 0$ and $i_j \neq i_k$ when $j \neq k$, then $\phi(a_1 a_2 \dots a_n) = 0$.*

Proof. We use induction w.r.t. n . For $n = 1$ the statement is obvious. Suppose that it holds for $l < n$. We have

$$\begin{aligned} \phi(a_1 a_2 \dots a_n) &= \phi(a_1^0 a_2 \dots a_n) + \phi(a_1) \phi(1_{i_1} a_2 \dots a_n) \\ &= \phi(a_1^0 a_2 \dots a_n) + \phi(a_1) \phi(a_2 \dots a_n). \end{aligned}$$

If $j \neq 1$, then $\phi(a_2 \dots a_n) = 0$ from the inductive assumption. If $j = 1$, then $\phi(a_1) = 0$, which gives

$$\phi(a_1 a_2 \dots a_n) = \phi(a_1^0 a_2 \dots a_n) = \phi(a_1^0 a_2^0 a_3 \dots a_n) + \phi(a_2) \phi(a_1^0 1_{i_2} a_3 \dots a_n)$$

As before, consider two cases. In the first case, namely $j = 2$ we have $\phi(a_2) = 0$, whereas in the second case, for $j \neq 2$ we have two possibilities:

1. $m \leq 1$. If $i_1 < i_2$, then $\phi(a_1^0 1_{i_2} a_3 \dots a_n) = \phi(a_1^0 a_3 \dots a_n) = 0$ from the inductive assumption. In turn, if $i_1 > i_2$, then $\phi(a_1^0 1_{i_2} a_3 \dots a_n) = 0$, which follows from Definition 3.1.
2. $m > 1$. Then $\phi(a_1^0 1_{i_2} a_3 \dots a_n) = \phi(a_1^0 a_3 \dots a_n) = 0$ from the inductive assumption.

Thus, in all considered cases, either $\phi(a_1^0 1_{i_2} a_3 \dots a_n) = 0$, or $\phi(a_2) = 0$, hence

$$\phi(a_1 a_2 \dots a_n) = \phi(a_1^0 a_2^0 a_3 \dots a_n).$$

By continuing this reasoning we obtain $\phi(a_1 a_2 \dots a_n) = \phi(a_1^0 a_2^0 \dots a_n^0)$, which is equal to zero by Definition 3.2. $\blacksquare$

LEMMA 6.2. *Suppose that $(i_1, \dots, i_n) \sim P$, where $i_1 \neq i_2 \neq \dots \neq i_n$ and where P is a crossing partition. Then the mixed moment $\phi(a_1 \dots a_n)$ is expressed in terms of products of at least $p + 1$ marginal moments, where p is the number of blocks of P .*

Proof. For $n = 4$, where the induction starts, the statement is true since $\phi(abab) = \phi(a^2)\phi^2(b)$ for $m = 1$ (monotone case) and

$$\phi(abab) = \phi^2(b)\phi(a^2) - \phi^2(b)\phi^2(a) + \phi^2(a)\phi(b^2)$$

for $m > 1$ (then, by Corollary 4.4, moments of order 4 agree with those in the free case). Let $n > 4$ and suppose that $i_1 \neq \dots \neq i_m < i_{m+1} < \dots < i_r > i_{r+1}$. In the formula

$$\phi(a_1 a_2 \dots a_n) = \phi(a_1^0 a_2 \dots a_n) + \phi(a_1)\phi(a_2 \dots a_n)$$

the second summand is a product of at least $p + 1$ factors since either $(i_2, \dots, i_n)$ corresponds to a non-crossing partition P' , and then P' must have p blocks, or $(i_2, \dots, i_n)$ corresponds to a crossing partition, in which case we use the inductive assumption. In turn, we can write

$$\phi(a_1^0 a_2 \dots a_n) = \phi(a_1^0 a_2^0 a_3 \dots a_n) + \phi(a_2)\phi(a_1^0 a_3 \dots a_n)$$

and use a similar argument to conclude that the second summand factorizes into at least $p + 1$ factors. Continuing this reasoning, we can reduce the proof to the case of $\phi(a_1^0 \dots a_r^0 a_{r+1} \dots a_n)$, which vanishes by Lemma 3.4. $\blacksquare$

LEMMA 6.3. *Suppose that $(i_1, \dots, i_n) \sim P \in \mathcal{ONC}_n(m)$, where $i_1 \neq i_2 \neq \dots \neq i_n$. Then the mixed moment $\phi(a_1 \dots a_n)$ is a product of exactly p marginal moments, where p is the number of blocks of P .*

Proof. We use induction w.r.t. n . For $n \leq 2m$, we get the same expression for the moments as in the free case by Corollary 4.4 and thus a product of p marginal moments [Sp]. Thus let $n > 2m$ and suppose that $i_1 \neq \dots \neq i_m < \dots < i_r > i_{r+1} \neq \dots \neq i_n$.

Since $(i_1, \dots, i_n)$ is non-crossing, hence there exists $1 \leq j \leq r$ such that $i_j \neq i_k$ for every $k \neq j$. Then

$$\begin{aligned}\phi(a_1 \dots a_n) &= \phi(a_1 \dots a_{j-1} a_j^0 a_{j+1} \dots a_n) + \phi(a_j) \phi(a_1 \dots a_{j-1} 1_{i_j} a_{j+1} \dots a_n) \\ &= \phi(a_j) \phi(a_1 \dots a_{j-1} a_{j+1} \dots a_n)\end{aligned}$$

by Lemma 3.5 since a_j^0 is a singleton of mean zero. To complete the proof, it is now enough to use the induction argument since $(i_1, \dots, i_{j-1}, i_{j+1}, \dots, i_n) \sim \mathcal{ONC}_{n-1}(m)$. $\blacksquare$

LEMMA 6.4. *Suppose that $i_1 \neq i_2 \neq \dots \neq i_n$ and that $(i_1, \dots, i_n) \sim P \in \mathcal{ONC}_n \setminus \mathcal{ONC}_n(m)$. Then the mixed moment $\phi(a_1 \dots a_n)$ is a product of at least $p+1$ marginal moments, where p is the number of blocks of P .*

Proof. Consider first the case when there exists $1 \leq r \leq n$ such that $i_1 \neq \dots \neq i_m < \dots < i_r > i_{r+1}$ and that for every $1 \leq j \leq r$ there exists $r+2 \leq k \leq n$ such that $i_j = i_k$. Then the second summand in the expression

$$\phi(a_1 \dots a_n) = \phi(a_1^0 a_2 \dots a_n) + \phi(a_1) \phi(a_2 \dots a_n)$$

is a product of at least $p+1$ factors since the partition associated with $(i_2, \dots, i_n)$ has p blocks. Hence it suffices to consider the first summand

$$\phi(a_1^0 a_2 \dots a_n) = \phi(a_1^0 a_2^0 a_3 \dots a_n) + \phi(a_2) \phi(a_1^0 a_3 \dots a_n)$$

and apply the same argument, using the property of m -monotone units. This reduces the proof to the case of $\phi(a_1^0 \dots a_r^0 a_{r+1} \dots a_n)$, which vanishes by Lemma 3.4. The general case can always be reduced to that considered above by means of Lemma 3.5 and Lemma 6.1. $\blacksquare$

7. CENTRAL LIMIT THEOREM

We will formulate now the central limit theorem for the normalized sums of m -monotone random variables. Then we shall give a recurrence relation for the sequence of their Cauchy transforms.

THEOREM 7.1. *Let $\{X_i\}_{i=1}^\infty$ be identically distributed m -monotone random variables with respect to ϕ of mean zero and variance one, i.e. $\phi(X_i) = 0$ and $\phi(X_i^2) = 1$. Then*

$$\lim_{N \rightarrow \infty} \phi \left(\left[\frac{X_1 + X_2 + \dots + X_N}{\sqrt{N}} \right]^n \right) = \begin{cases} 0 & \text{if } n = 2k+1 \\ \frac{|\mathcal{ONC}_{2k}^2(m)|}{k!} & \text{if } n = 2k \end{cases}.$$

Proof. Since mixed moments are invariant under order-preserving injections, we get

$$\begin{aligned}\phi((X_1 + \dots + X_N)^n) &= \sum_{1 \leq k_1, \dots, k_n \leq N} \phi(X_{k_1} \dots X_{k_n}) \\ &= \sum_{P \in \mathcal{OP}_n} \binom{N}{b(P)} m(P)\end{aligned}$$

where $\mathcal{OP}_n$ denotes all ordered partitions of the set $\{1, \dots, n\}$,

$$m(P) = \phi(X_{k_1} \dots X_{k_n}) \text{ for } (k_1, \dots, k_n) \sim P$$

and $b(P)$ denotes the number of blocks of P .

Let us now analyze the contribution from every ordered partition P for large N . If there is a singleton in P , then by Lemma 6.1, $m(P) = 0$. Thus suppose that P has no singletons. If $n = 2k + 1$, then $b(P) \leq k$ and after dividing $\binom{N}{b(P)}$ by $N^{k+1/2}$ we get a contribution of order smaller or equal to $N^{-1/2}$. If $n = 2k$, then either $b(P) < k$, when the contribution is of order smaller or equal to N^{-1} , or $b(P) = k$, when we get $1/b(P)!m(P)$, but only for those ordered pair-partitions for which $m(P)$ factorizes into k factors. Lemmas 6.2-6.4 imply that if $P \in \mathcal{ONC}_{2k}^2(m)$, then $m(P)$ factorizes into $b(P)$ factors, whereas if $P \notin \mathcal{OP}_{2k}^2 \setminus \mathcal{ONC}_{2k}^2(m)$, then $m(P)$ factorizes into more than $b(P)$ factors. This implies the latter do not contribute to the limit. In turn, the contribution from the former is $1/k!$ since variance is assumed to be equal to one. This completes the proof. $\blacksquare$

Remark. If $m = 1$, then using Theorem 7.1, we get in the CLT

$$M_n^{(1)} = \begin{cases} 0 & \text{if } n = 2k + 1 \\ \frac{1}{2^k} \binom{2k}{k} & \text{if } n = 2k \end{cases}$$

the moments of the arcsine law supported on $[-\sqrt{2}, \sqrt{2}]$.

THEOREM 7.2. *The Cauchy transforms $G^{(m)}(z)$ of the limit laws obtained in the CLT for the monotone hierarchy satisfy the recurrence*

$$G^{(1)}(z) = \frac{1}{\sqrt{z^2 - 2}} \quad , \quad G^{(m)}(z) = \frac{1}{z - G^{(m-1)}(z)} \quad \text{for } m \geq 2$$

for $z \in \mathbb{C}_+ = \{z : \Im z > 0\}$.

Proof. That $G^{(1)}(z)$ is the Cauchy transform of the arcsine law is well-known (see, for instance [M2]). Thus let $m \geq 2$. Denote by $N_{2n}(m)$ the cardinality of $\mathcal{ONC}_{2n}^2(m)$. We will express $N_{2n+2}(m)$ in terms of $N_{2k}(m)$ for $k \leq n$. From the set $\{2, 4, \dots, 2n + 2\}$ choose an even number which will form a pair with 1 and denote it $2k$. Using the fact that partitions from $\mathcal{ONC}_{2n+2}^2(m)$ are non-crossing, we can separate blocks which are inner w.r.t. $\{1, 2k\}$ from other blocks (subsets of $\{2k + 1, \dots, 2n + 2\}$). The collection

of the latter forms a partition of $\{2k+1, \dots, 2n+2\}$ associated with $\mathcal{ONC}_{2n+2-2k}^2(m)$ and there are exactly $\binom{n+1}{n+1-k} N_{2n+2-2k}(m)$ such partitions. On the other hand, blocks which are inner w.r.t. $\{1, 2k\}$ form a partition associated with $\mathcal{ONC}_{2k-2}^2(m-1)$, and there are $\binom{k}{k-1} N_{2k-2}(m-1)$ ways of choosing them. Thus

$$N_{2n+2}(m) = \sum_{k=1}^{n+1} k \binom{n+1}{k} N_{2k-2}(m-1) N_{2n-2k+2}(m).$$

Dividing both sides of the equation by $(n+1)!$ and denoting the moments of the central limit laws by $M_n^{(m)}$, we obtain

$$\begin{aligned} \frac{N_{2n+2}(m)}{(n+1)!} &= \sum_{k=1}^{n+1} \frac{N_{2k-2}(m-1)}{(k-1)!} \frac{N_{2n-2k+2}(m)}{(n-k+1)!} \\ M_{2n+2}^{(m)} &= \sum_{k=1}^{n+1} M_{2k-2}^{(m-1)} M_{2n-2k+2}^{(m)}. \end{aligned}$$

This identity allows us to derive a recurrence for the Cauchy transforms:

$$\begin{aligned} G^{(m)}(z) &= \sum_{n=0}^{\infty} M_{2n}^{(m)} z^{-2n-1} = \frac{1}{z} + \sum_{n=0}^{\infty} M_{2n+2}^{(m)} z^{-2n-3} \\ &= \frac{1}{z} + \frac{1}{z} \sum_{n=0}^{\infty} \sum_{k=1}^{n+1} M_{2k-2}^{(m-1)} z^{-2k+1} M_{2n-2k+2}^{(m)} z^{-2n+2k-3} \\ &= \frac{1}{z} + \frac{1}{z} \left(\sum_{n=0}^{\infty} M_{2n}^{(m)} z^{-2n-1} \right) \left(\sum_{n=0}^{\infty} M_{2n}^{(m-1)} z^{-2n-1} \right) \\ &= \frac{1}{z} + \frac{G^{(m)}(z) G^{(m-1)}(z)}{z}, \end{aligned}$$

which leads to

$$G^{(m)}(z) = \frac{1}{z - G^{(m-1)}(z)},$$

and this completes the proof. ■

The table given below gives even moments of lowest orders obtained in the CLT for the monotone hierarchy as well as those of the Wigner law which corresponds to $m = \infty$. Let us observe that for given m , moments of orders smaller or equal to $2m$ agree with the moments of the Wigner law.

	$n = 2$	$n = 4$	$n = 6$	$n = 8$	$n = 10$
$m = 1$	1	$3/2$	$5/2$	$35/8$	$63/8$
$m = 2$	1	2	$9/2$	$21/2$	$199/8$
$m = 3$	1	2	5	$27/2$	$75/2$
$m = 4$	1	2	5	14	$83/2$
$m = \infty$	1	2	5	14	42

Table 1. Low-order moments of the m -monotone central limit laws.

Example 7.1. If $m = 2$, then the Cauchy transform of the central limit law takes the form

$$G^{(2)}(z) = \frac{1}{z - \frac{1}{z\sqrt{1 - \frac{2}{z^2}}}}.$$

In order to compute the absolutely continuous part of the limit measure for $m = 2$ we use the Stieltjes inversion formula:

$$\begin{aligned} f^{(2)}(x) &= -\frac{1}{\pi} \lim_{y \rightarrow 0^+} \Im G^{(2)}(x + iy) \\ &= \begin{cases} \frac{1}{\pi} \frac{\sqrt{2 - x^2}}{1 + x^2(2 - x^2)} & \text{for } |x| \leq \sqrt{2} \\ 0 & \text{for } |x| > \sqrt{2} \end{cases}. \end{aligned}$$

The singular part of the limit measure is obtained by computing the residua at the real poles of $G^{(2)}(z)$, namely

$$\text{Res}_{\sqrt{\sqrt{2}+1}} G^{(2)}(z) = \text{Res}_{-\sqrt{\sqrt{2}+1}} G^{(2)}(z) = \frac{2 - \sqrt{2}}{4}.$$

Figure 3 below shows both the absolutely continuous and the singular parts of the central limit law for $m = 2$.

Using similar computations and Mathematica packet for $m = 3$, we obtain a measure with the absolutely continuous part

$$f^{(3)}(x) = \begin{cases} \frac{1}{\pi} \frac{\sqrt{2 - x^2}}{x^2 + (x^2 - 1)^2(2 - x^2)} & \text{for } |x| \leq \sqrt{2} \\ 0 & \text{for } |x| > \sqrt{2} \end{cases}.$$

and the singular part consisting of two atoms of masses approximately equal to 0,099 each, concentrated at $\pm 1,685$, see Figure 4. An analogous picture for $m = 4$ is given in Figure 5.




Figure 3. The central limit law for $m = 2$.




Figure 4. The central limit law for $m = 3$.




Figure 5. The central limit law for $m = 4$.

8. POISSON'S LIMIT THEOREM.

In this Section we prove the Poisson's limit theorem.

THEOREM 8.1. *Let $m \in \mathbb{N}$ and suppose that for every $N \in \mathbb{N}$ the variables $X_{1,N}, \dots, X_{N,N}$ are identically distributed and m -monotone with respect to ϕ_N . If $N\phi_N(X_{i,N}^k) \rightarrow \lambda^k$ for every natural k , where $\lambda > 0$, then*

$$\lim_{N \rightarrow \infty} \phi_N((X_{1,N} + \dots + X_{N,N})^n) = \sum_{q=1}^n \frac{\lambda^q}{q!} |\mathcal{ONC}_n(q, m)|$$

where $\mathcal{ONC}_n(q, m)$ denotes the subset of $\mathcal{ONC}_n(m)$ consisting of partitions which have q blocks.

Proof. Using invariance of mixed moments under order-preserving injections, we get

$$\begin{aligned} \phi_N((X_{1,N} + \dots + X_{N,N})^n) &= \sum_{1 \leq k_1, \dots, k_n \leq N} \phi_N(X_{k_1,N} \dots X_{k_n,N}) \\ &= \sum_{P \in \mathcal{CP}_n} \binom{N}{b(P)} m_N(P) \end{aligned}$$

where $\mathcal{OP}_n$ denotes all ordered partitions of the set $\{1, \dots, n\}$,

$$m_N(P) = \phi_N(X_{k_1, N} \dots X_{k_n, N}) \text{ whenever } (k_1, \dots, k_n) \sim P$$

and $b(P)$ denotes the number of blocks of P .

Lemmas 6.2-6.4 imply that if $P \in \mathcal{ONC}_n(m)$, then $m_N(P)$ factorizes into $b(P)$ factors, whereas if $P \notin \mathcal{ONC}_n(m)$, then $m(P)$ factorizes into more than $b(P)$ factors. This implies that the latter do not contribute to the limit. In turn, the contribution from $P \in \mathcal{ONC}_n(m)$, depending on $b(P)$, gives the desired formula. $\blacksquare$

Again, as in the case of the central limit theorem, the recurrence for Cauchy transforms is the same as in the m -free case [L-F] and only the initial condition is different. We use the notations:

$$M_n^{(m)}(\lambda, q) = \frac{\lambda^q}{q!} |\mathcal{ONC}_n(q, m)| \quad , \quad H^{(m)}(\lambda, z) = \sum_{n, q=0}^{\infty} M_n^{(m)}(\lambda, q) z^{-n-1}$$


with $M_n^{(m)}(\lambda, 0) = \delta_{n,0}$.

LEMMA 8.2. *The hierarchy of generating functions $\{H^{(m)}\}_{m=0}^{\infty}$ satisfies the recurrence relation*

$$H^{(m)}(\lambda, z) = \frac{1 - H^{(m-1)}(\lambda, z)}{z - zH^{(m-1)}(\lambda, z) - \lambda}$$

for $m = 2, 3, \dots$ and $H^{(1)}(\lambda, z)$ is the product log function described in [M2].

Proof. The case $m = 1$ follows from [M2]. Suppose that $m \geq 2$. To get an ordered non-crossing partition of $\{1, \dots, n\}$ we pick the elements that will be put in the same block as the first element; denote this block by $\{1, p_1, \dots, p_{r-1}\}$ as in the diagram given below, with $Q_1, Q_2, \dots, Q_r$ denoting subpartitions of the intervals $(1, p_1 - 1)$, $(p_1 + 1, \dots, p_2 - 1), \dots, (p_{r-1} + 1, n]$, respectively.



As compared with the m -free case we need to take into account the number of possible block colorings. Denoting $N_n(q, m) = |\mathcal{ONC}_n(q, m)|$, we get

$$\begin{aligned} N_n(q, m) &= \sum_{r=1}^n \sum_{k_1 + \dots + k_r = n-1} \sum_{q_1 + \dots + q_r = q-1} \frac{q!}{q_1! q_2! \dots q_r!} \\ &\times N_{k_1-1}(q_1, m-1) \dots N_{k_{r-1}-1}(q_{r-1}, m-1) N_{k_r}(q_r, m) \end{aligned}$$

for $n, k_1, \dots, k_{r-1} \geq 1$; $k_r, b_1, \dots, b_r \geq 0$, where

$$k_1 = p_1 - 1, k_2 = p_2 - p_1, \dots, k_{r-1} = p_{r-1} - p_{r-2}$$

since there are $q!/(q_1! \dots q_r!)$ ways to choose *sets* of colors for subpartitions $Q_1, \dots, Q_r$ from among $q = q_1 + \dots + q_r$ colors and then ‘admissible’ colorings of these sets give numbers $N_{k_1-1}(q_1, m-1), \dots, N_{k_r}(q_r, m)$. Now, multiplying both sides by $\lambda^q/q!$, we get

$$\begin{aligned} M_n^{(m)}(\lambda, q) &= \lambda \sum_{r=1}^n \sum_{k_1+\dots+k_r=n-1} \sum_{q_1+\dots+q_r=q-1} \\ &\quad M_{k_1-1}^{(m-1)}(\lambda, q_1) \times \dots M_{k_{r-1}-1}^{(m-1)}(\lambda, q_{r-1}) M_{k_r}^{(m)}(\lambda, q_r) \end{aligned}$$

Then we have

$$\begin{aligned} H^{(m)}(\lambda, z) &= \sum_{n,q=0}^{\infty} M_n^{(m)}(\lambda, q) z^{-n-1} \\ &= \frac{1}{z} + \frac{\lambda}{z} \sum_{n,q=1}^{\infty} \sum_{r=1}^n \sum_{k_1+\dots+k_r=n-1} \sum_{q_1+\dots+q_r=q-1} M_{k_1-1}^{(m-1)}(\lambda, q_1) z^{-k_1} \\ &\quad \times \dots M_{k_{r-1}-1}^{(m-1)}(\lambda, q_{r-1}) z^{-k_{r-1}} M_{k_r}^{(m)}(\lambda, q_r) z^{-k_r-1} \\ &= \frac{1}{z} + \frac{\lambda}{z} \sum_{r=1}^{\infty} \left(\sum_{\beta, \nu=0}^{\infty} M_{\nu}^{(m-1)}(\lambda, \beta) z^{-\nu-1} \right)^{r-1} \sum_{\mu, \alpha=0}^{\infty} M_{\mu}^{(m)}(\lambda, \alpha) z^{-\mu-1} \\ &= \frac{1}{z} + \frac{\lambda H^{(m)}(\lambda, z)}{z(1 - H^{(m-1)}(\lambda, z))}, \end{aligned}$$

and therefore

$$H^{(m)}(\lambda, z) = \frac{1 - H^{(m-1)}(\lambda, z)}{z - zH^{(m-1)}(\lambda, z) - \lambda}$$

which completes the proof. ■

9. INVARIANCE PRINCIPLE

Let $(X_i)_{i \in \mathbb{N}} \in \mathcal{A}$ be a sequence of identically distributed random variables of mean zero and variance one, which are m -monotone with respect to a state ϕ on $\mathcal{A}$. Below we shall investigate the asymptotic behavior of normalized sums

$$S_N(f) = \frac{1}{\sqrt{N}} \sum_{i=[sN]+1}^{[tN]} X_i \tag{9.1}$$

for $f = \chi(s, t] \in \Theta$ (see (5.1)) as $N \rightarrow \infty$.

It is convenient to give a formula for mixed moments of sums indexed by functions which have identical or disjoint supports.

THEOREM 9.1. *Suppose $f_1, f_2, \dots, f_n \in \Theta$ have identical or disjoint supports and let $\sigma(f) = \{\sigma_1, \sigma_2, \dots, \sigma_p\}$ be the partition defined by the tuple $(f_1, f_2, \dots, f_n)$, i.e. each block σ_k consists of all those indices j , for which f_j coincide. Then*

$$\lim_{N \rightarrow \infty} \phi(S_N(f_1)S_N(f_2) \dots S_N(f_n)) = \frac{1}{b_1!b_2! \dots b_p!} \sum_{\substack{P \in \mathcal{ONC}_n^2(m) \\ P \sim (f_1, f_2, \dots, f_n)}} \prod_{\{\alpha, \beta\} \in P} \langle f_\alpha, f_\beta \rangle \quad (9.2)$$

where $b_k = |\sigma_k|/2$ for $k = 1, \dots, p$, with the understanding that $\mathcal{ONC}_n^2(m) = \emptyset$ for n odd.

Proof. It is clear that if n is odd, we get zero contribution - we understand here that in that case $\mathcal{ONC}_n^2(m) = \emptyset$. Thus, let $n = 2k$ and let $f_k = \chi_{(s_k, t_k]}$ for $k = 1, \dots, n$. We have

$$\begin{aligned} \phi(S_N(f_1)S_N(f_2) \dots S_N(f_n)) &= \frac{1}{N^{n/2}} \sum_{i_1=[s_1N]+1}^{[t_1N]} \dots \sum_{i_n=[s_nN]+1}^{[t_nN]} \phi(X_{i_1} \dots X_{i_n}) \\ &= \frac{1}{N^{n/2}} \sum_{P \in \mathcal{ONC}_n^2(m)} A_P(f_1, f_2, \dots, f_n; N) + \mathcal{O}(1/N) \end{aligned}$$

where $A_P(f_1, f_2, \dots, f_n; N)$ denotes the number of tuples $(i_1, i_2, \dots, i_n)$ such that

$$[s_jN] < i_j \leq [t_jN] \text{ for } j = 1, \dots, n$$

and $(i_1, i_2, \dots, i_n) \sim P$. Now, Lemma 6.1 together with the mean zero assumption result in the fact that only contributions from pair partitions are of order 1 - other partitions must have fewer than k blocks and thus their contribution is at most of order $1/N$ as in the CLT. Moreover, moments associated with ordered pair partitions which are not in $\mathcal{ONC}_n^2(m)$ factorize into more than k factors by Lemmas 6.2-6.4 and thus vanish by the mean zero assumption. Finally, if $P \in \mathcal{ONC}_n^2(m)$ and $P \sim (f_1, f_2, \dots, f_n)$, then

$$A_P(f_1, f_2, \dots, f_n; N) = \binom{N_1}{b_1} \binom{N_2}{b_2} \dots \binom{N_p}{b_p}$$

where $N_j = [t^{(j)}N] - [s^{(j)}N]$ and $t^{(j)}, s^{(j)}$ denote the end-points of f_r for any $r \in \sigma_j$, where $j = 1, \dots, p$ (thus N_j counts the number of possible values of those indices in the tuple $(i_1, i_2, \dots, i_n)$ which are associated with block σ_j). For such P we get

$$\begin{aligned} \frac{A_P(f_1, f_2, \dots, f_n)}{N^k} &= \prod_{j=1}^p \frac{N_j(N_j-1) \dots (N_j-b_j+1)}{b_j!N^{b_j}} \\ &\rightarrow \prod_{j=1}^p \frac{(t^{(j)} - s^{(j)})^{b_j}}{b_j!} \\ &= \frac{1}{b_1!b_2! \dots b_p!} \prod_{\{\alpha, \beta\} \in P} \langle f_\alpha, f_\beta \rangle \end{aligned}$$

whereas if $P \not\sim (f_1, f_2, \dots, f_n)$, then $A_P(f_1, f_2, \dots, f_n; N) = 0$, which finishes the proof. $\blacksquare$

10. m -MONOTONE GAUSSIAN OPERATORS

In this section we introduce the m -monotone Fock space as an m -th order iteration of the free Fock space, on which we give a realization of the limit moments in the invariance principle for m -monotone independence.

The m -free Fock space over a Hilbert space $\mathcal{H}$ [FL] is defined by the direct sum

$$\mathcal{F}_{(m)}(\mathcal{H}) = \mathbb{C}\Omega \oplus \bigoplus_{n=1}^m \mathcal{H}^{\otimes n}$$

with the canonical scalar product inherited from the free Fock space $\mathcal{F}(\mathcal{H})$, where $m \in \mathbb{N}$ and Ω is a unit vector called vacuum.

Let $\mathcal{H} = L^2(\mathbb{R}_+)$, then $\mathcal{H}^{\otimes n} \cong L^2(\mathbb{R}_+^n)$ and thus another direct sum

$$\mathcal{F}^{(m)}(\mathcal{H}) = \mathcal{F}_{(m)}(\mathcal{H}) \oplus \bigoplus_{n=m+1}^{\infty} L^2(\Delta_n^{(m)}) \quad (10.1)$$

can be viewed as a subspace of $\mathcal{F}(\mathbb{R}_+)$ with ‘order starting from the m -th level’, where

$$\Delta_n^{(m)} = \{(x_1, x_2, \dots, x_n) : x_1 > \dots > x_{n-m+1} > 0\}$$

for every $n \in \mathbb{N}$. We equip $\mathcal{F}^{(m)}(\mathcal{H})$ with the canonical scalar product inherited from $\mathcal{F}(\mathcal{H})$ and call it the m -monotone Fock space. In particular, $\mathcal{F}^{(1)}(\mathcal{H})$ is the monotone Fock space over $\mathcal{H}$ [M2, Lu].

If we denote by $\mathcal{P}^{(m)}$ the canonical projection from $\mathcal{F}(\mathcal{H})$ onto $\mathcal{F}^{(m)}(\mathcal{H})$, then $\mathcal{F}^{(m)}(\mathcal{H})$ is spanned by Ω and vectors of the form

$$f_1 \otimes_m \dots \otimes_m f_n := \mathcal{P}^{(m)}(f_1 \otimes \dots \otimes f_n)$$

where $f_1, \dots, f_n \in \mathcal{H}$ and $n \in \mathbb{N}$.

Define m -monotone creation operators on $\mathcal{F}^{(m)}(\mathcal{H})$ as bounded extensions of

$$\begin{aligned} a^{(m)}(f)\Omega &= f \\ a^{(m)}(f)(f_1 \otimes_m \dots \otimes_m f_n) &= f \otimes_m f_1 \otimes_m \dots \otimes_m f_n \end{aligned}$$

and the m -monotone annihilation operators $a^{(m)*}(f)$ as their adjoints.

It is easy to see that $a^{(m)*}(f)\Omega = 0$ and that

$$a^{(m)*}(f)(f_1 \otimes_m \dots \otimes_m f_n) = \begin{cases} \langle f_1, f \rangle f_2 \otimes_m \dots \otimes_m f_n & 1 \leq n \leq m \\ M_\psi f_2 \otimes_m \dots \otimes_m f_n & n > m \end{cases}$$

where $(M_{\psi}g)(x) = \psi(x)g(x)$ is the multiplication operator by function ψ and

$$\psi(x) = \langle f_1, f \rangle_x = \int_{y>x} f_1(y)\bar{f}(y)dy \quad (10.2)$$

(dependence of ψ on f and f_1 is suppressed). Note that if $\text{supp}f_1 \cap \text{supp}f > \text{supp}f_2$, then we have $\psi(x) = \langle f_1, f \rangle$ for every $x \in \text{supp}f_2$.

Now let us define vacuum states on all bounded operators T on the Fock spaces $\mathcal{F}^{(m)}(\mathcal{H})$:

$$\varphi(T) = \langle T\Omega, \Omega \rangle$$

(by abuse of notation we use the same notation for all m), and position operators

$$\omega^{(m)}(f) = a^{(m)}(f) + a^{(m)*}(f)$$

for $f \in \Theta$ given by (5.1), which may be called *m-monotone Gaussian operators*.

Our goal is now to express the limit moments obtained in the invariance principle in terms of the vacuum expectations of the *m-monotone Gaussian operators*. For clarity of exposition, we first consider the monotone case $m = 1$. A generalization to all natural m is rather straightforward. The combinatorial formulas for mixed moments even in the case of monotone Gaussians (with arcsine distribution) are new.

To every $\pi \in \mathcal{NC}_{2k}^2$ and $(f_1, f_2, \dots, f_{2k})$ we associate the operator

$$a_{\pi}(f_1, f_2, \dots, f_{2k}) = a^{\epsilon_1}(f_1)a^{\epsilon_2}(f_2) \dots a^{\epsilon_{2k}}(f_{2k}) \quad (10.3)$$

where $\epsilon_p = *$ and $\epsilon_q = 1$ whenever $\{p, q\}$ is a block of π . We also use notations (5.2) and (5.3).

LEMMA 10.1. *Let $\pi = \{\pi_1, \pi_2, \dots, \pi_k\} \in \mathcal{NC}_{2k}^2$ and $\pi \sim (f_1, f_2, \dots, f_{2k})$, where $f_1, f_2, \dots, f_n \in \Theta$ have identical or disjoint supports. Then*

$$\varphi(a_{\pi}(f_1, f_2, \dots, f_{2k})) = \psi_{\pi_1}(s^{(1)})\psi_{\pi_2}(s^{(2)}) \dots \psi_{\pi_k}(s^{(k)}) \quad (10.4)$$

where $\psi_{\pi_i}(x) = (\text{Inn}(\pi_i) + 1)^{-1}(t^{(i)} - x)$ is the linear function associated with block π_i for $x \in [s^{(i)}, t^{(i)}]$, $i = 1, \dots, k$.

Proof. Clearly, if $k = 1$ and $\pi = \{\{1, 2\}\}$, then

$$a_{\pi}\Omega = a^*(f_1)a(f_1)\Omega = (t^{(1)} - s^{(1)})\Omega = \psi_{\pi_1}(s^{(1)})\Omega,$$

which gives the formula for $k = 1$.

Claim: Abbreviating notation (10.3) to a_{π} , we get

$$a_{\pi}(g_1 \otimes_1 \dots \otimes_1 g_n) = \prod_{\substack{1 \leq i \leq k \\ \text{supp}f_i \neq \text{supp}g_1}} \psi_{\pi_i}(s^{(i)}) \left(\prod_{\substack{1 \leq j \leq k \\ \text{supp}f_j = \text{supp}g_1}} M_{\psi_j}g_1 \right) \otimes_1 \dots \otimes_1 g_n$$

for $g_1, \dots, g_n \in \mathcal{H}$ with $\text{supp}g_1 = \text{supp}f_{2k}$ or $\text{supp}g_1 < \text{supp}f_{2k}$, where $n \in \mathbb{N}$.

Suppose this formula holds for $\pi \in \mathcal{NC}_{2r}^2$ for $1 \leq r \leq k-1$. If $\pi = \pi' \cup \pi''$, where π' is a non-crossing partition of $\{1, \dots, 2r\}$ and π'' is a non-crossing partition of $\{2r+1, \dots, 2k\}$, then it is obvious that the claim holds for π . Therefore, suppose that $\pi = \pi' \cup \{1, 2k\}$. Then, using the inductive assumption (and abbreviated self-explanatory notation $a_{\pi'}$), we get

$$\begin{aligned} a_{\pi}(g_1 \otimes_1 \dots \otimes_1 g_n) &= a^*(f_1) a_{\pi'}(f_{2k} \otimes_1 g_1 \otimes_1 \dots \otimes_1 g_n) \\ &= \prod_{\substack{2 \leq j \leq k \\ f_j \neq f_{2k}}} \psi_{\pi_i}(s^{(i)}) a^*(f_1) \left(\prod_{\substack{2 \leq j \leq k \\ f_j = f_{2k}}} M_{\psi_j} f_{2k} \otimes_1 g_1 \otimes_1 \dots \otimes_1 g_n \right) \\ &= \prod_{\substack{2 \leq j \leq k \\ f_i \neq f_{2k}}} \psi_{\pi_i}(s^{(i)}) (M_{\psi} g_1 \otimes_1 \dots \otimes_1 g_n), \end{aligned}$$

where

$$\psi(x) = \begin{cases} \psi_{\pi_1}(x) & \text{if } \text{supp} g = \text{supp} f_{2k} \\ \psi_{\pi_1}(s^{(1)}) & \text{if } \text{supp} g < \text{supp} f_{2k} \end{cases}$$

follows from

$$\psi(x) = \int_x^{t^{(1)}} \prod_{\substack{2 \leq j \leq k \\ f_j = f_{2k}}} M_{\psi_j} 1 dx = \int_x^{t^{(1)}} (t^{(1)} - y)^{b_1-1} dy = \frac{1}{b_1} (t^{(1)} - x)^{b_1}$$

where $b_1 = \text{Inn}(\pi_1) + 1$, whenever $\text{supp} g = \text{supp} f_{2k}$, whereas if $\text{supp} g \neq \text{supp} f_{2k}$, then $\psi(x) = \psi_{\pi_1}(s^{(1)})$. This proves the claim, from which (10.8) follows easily. $\blacksquare$

Example 10.1. Let us evaluate the moment $\langle a_{\pi} \Omega, \Omega \rangle$, where π is given by the diagram in Figure 2 (see Section 5). In view of Lemma 10.1, we get

$$\langle a_{\pi} \Omega, \Omega \rangle = \frac{1}{2} t^2 (t' - t)^2$$

since $\phi_1(x) = 1/2(t - x)$, $\phi_2(x) = t' - x$, $\phi_3(x) = t - x$, $\phi_4(x) = t' - x$ (cf. Example 5.3).

For comparison, note that in the case of vacuum expectations of free creation and annihilation operators there is a multiplicative formula similar to (10.4), except that on the RHS we get $(t^{(1)} - s^{(1)}) \dots (t^{(k)} - s^{(k)})$. Thus, the difference is that in the monotone case the product encodes additional information about inner blocks.

Moreover, as the lemma below demonstrates, the number of admissible colorings of π can also be expressed in terms of the same information about inner blocks.

LEMMA 10.2. *Let $\pi = \{\pi_1, \pi_2, \dots, \pi_k\} \in \mathcal{NC}_{2k}^2$ and $\pi \sim (f_1, f_2, \dots, f_{2k})$, where $f_1, f_2, \dots, f_n \in \Theta$ have identical or disjoint supports. Then*

$$\frac{c_{\pi}(f_1, f_2, \dots, f_{2k})}{b_1! b_2! \dots b_p!} = \prod_{i=1}^k (\text{Inn}(\pi_i) + 1)^{-1} \quad (10.5)$$

with the same notation as in Theorem 9.1.

Proof. Clearly, the formula is true for $k = 1$ and $\pi = \{1, 2\}$. Suppose the formula holds for non-crossing pair partitions of sets consisting of $2k - 2$ elements and let $\pi \in \mathcal{NC}_{2k}^2$.

Case 1. Suppose $\pi = \pi' \cup \pi''$, where $\pi' = \{\pi'_1, \dots, \pi'_r\}$ is a non-crossing pair partition of $\{1, \dots, 2r\}$ and $\pi'' = \{\pi''_1, \dots, \pi''_{k-r}\}$ is a non-crossing pair-partition of $\{2r+1, \dots, 2k\}$. Let a_i denote the number of non-empty blocks of the partition $\tau^{(i)} = \{\pi'_1 \cap \sigma_i, \dots, \pi'_r \cap \sigma_i\}$, where $i = 1, \dots, p$. Since $\pi \sim (f_1, \dots, f_{2k})$, there are $\binom{b_i}{a_i}$ ways to choose colors for blocks of $\tau^{(i)}$. Since colors are chosen independently for different supports, we get

$$\begin{aligned} \frac{c_\pi(f_1, \dots, f_{2k})}{b_1! \dots b_p!} &= \binom{b_1}{a_1} \dots \binom{b_p}{a_p} \frac{c_{\pi'}(f_1, \dots, f_{2r}) c_{\pi''}(f_{2r+1}, \dots, f_{2k})}{b_1! \dots b_p!} \\ &= \frac{c_{\pi'}(f_1, \dots, f_{2r})}{a_1! \dots a_p!} \frac{c_{\pi''}(f_{2r+1}, \dots, f_{2k})}{(b_1 - a_1)! \dots (b_p - a_p)!} \\ &= \prod_{i=1}^r (\text{Inn}(\pi'_i) + 1)^{-1} \prod_{j=1}^{k-r} (\text{Inn}(\pi''_j) + 1)^{-1} \\ &= \prod_{i=1}^k (\text{Inn}(\pi_i) + 1)^{-1} \end{aligned}$$

which gives (10.9) for the partition π .

Case 2. Suppose $\pi = \pi' \cup \{1, 2k\}$, where $\pi' = \{\pi_2, \dots, \pi_k\}$ is a non-crossing pair-partition of the set $\{2, \dots, 2k-1\}$. Thus all blocks of π' are inner w.r.t. the block $\pi_1 = \{1, 2k\}$ and this implies that π_1 must be colored by 1. Therefore, there are exactly the same numbers of ways of coloring π as well as π' . Choosing b_1 to be the number of blocks of π with the same support as $f_1 = f_{2k}$, we get

$$\begin{aligned} c_\pi(f_1, \dots, f_{2k}) &= c_{\pi'}(f_2, \dots, f_{2k-1}) \\ &= (b_1 - 1)! b_2! \dots b_p! \prod_{i=2}^k (\text{Inn}(\pi_i) + 1)^{-1} \\ &= b_1! b_2! \dots b_p! \prod_{i=1}^k (\text{Inn}(\pi_i) + 1)^{-1} \end{aligned}$$

since $b_1 = \text{Inn}(\pi_1) + 1$, which gives (10.9) for the partition π . This completes the proof. $\blacksquare$

THEOREM 10.3. *Suppose $f_1, f_2, \dots, f_n \in \Theta$ have identical or disjoint supports. Then*

$$\varphi(\omega^{(1)}(f_1) \omega^{(1)}(f_2) \dots \omega^{(1)}(f_n)) = \frac{1}{b_1! b_2! \dots b_p!} \sum_{\substack{P \in \mathcal{MNC}_n^2 \\ P \sim (f_1, f_2, \dots, f_n)}} \prod_{\{\alpha, \beta\} \in P} \langle f_\alpha, f_\beta \rangle$$

with the same notation as in Theorem 9.2.

Proof. If $n = 2k$, we have

$$\begin{aligned}
\varphi(\omega^{(1)}(f_1) \dots \omega^{(1)}(f_{2k})) &= \sum_{\substack{\pi \in \mathcal{NC}_{2k}^2 \\ \pi \sim (f_1, \dots, f_n)}} \langle a_\pi(f_1, \dots, f_{2k}) \Omega, \Omega \rangle \\
&= \sum_{\substack{\pi \in \mathcal{NC}_{2k}^2 \\ \pi \sim (f_1, \dots, f_n)}} \psi_{\pi_1}(s^{(1)}) \dots \psi_{\pi_k}(s^{(k)}) \\
&= \sum_{\substack{\pi \in \mathcal{NC}_{2k}^2 \\ \pi \sim (f_1, \dots, f_n)}} \prod_{i=1}^k \frac{t^{(i)} - s^{(i)}}{\text{Inn}(\pi_i) + 1} \\
&= \frac{1}{b_1! \dots b_p!} \sum_{\substack{\pi \in \mathcal{NC}_{2k}^2 \\ \pi \sim (f_1, \dots, f_n)}} c_\pi(f_1, \dots, f_{2k}) \prod_{\{\alpha, \beta\} \in \pi} \langle f_\alpha, f_\beta \rangle \\
&= \frac{1}{b_1! \dots b_p!} \sum_{\substack{P \in \mathcal{MNC}_{2k}^2 \\ P \sim (f_1, \dots, f_n)}} \prod_{\{\alpha, \beta\} \in P} \langle f_\alpha, f_\beta \rangle
\end{aligned}$$

where we used Lemmas 10.2-10.3. It is clear that if n is odd, then the expectation vanishes. This completes the proof. $\blacksquare$

In particular, the combinatorics of inner blocks allows us to express the moments of the arcsine law as a sum $M_{2k}^{(1)} = \sum_{\pi \in \mathcal{NC}_{2k}^2} \varphi(a_\pi)$, where the vacuum expectations $\varphi(a_\pi)$ are multiplicative functions over blocks of π , namely

$$\varphi(a_\pi) = \psi_{\pi_1}(1) \psi_{\pi_2}(1) \dots \psi_{\pi_k}(1).$$

Note that in the combinatorics of block depths given in [AB], the vacuum expectations $\varphi(b_\pi)$ on the algebra generated by creation and annihilation operators on the interacting Fock space do not have the above property, i.e.

$$\varphi(b_\pi) \neq \beta_{d(1)} \beta_{d(2)} \dots \beta_{d(k)}$$

with β_j 's as in (1.5) and $d(j)$'s denoting the depths of π_j 's, although the moments of the arcsine law are equal to the sums of such products.

Finally, let us remark that Lemmas 10.1-10.2 and Theorem 10.3 can be generalized to arbitrary $m \in \mathbb{N}$ by setting $\text{Inn}(\pi_k) = 0$ for blocks of depth $d_k < m$ and keeping $\text{Inn}(\pi_j)$ as in (5.2) for blocks of depth $d_j \geq m$. This means that up to depth m we have the combinatorics of free probability and starting from depth m we have the combinatorics of monotone probability. Proofs are very similar but slightly more technical and are omitted.

BIBLIOGRAPHY

- [AB] L. Accardi, M. Bożejko, Interacting Fock spaces and Gaussianization of probability measures, *Inf. Dim. Anal. Quant. Prob. Rel. Topics* **1** (1998), 663-670.
- [Av] D. Avitzour, Free products of C^* -algebras, *Trans. Amer. Math. Soc.* **271** (1982), 423-465.
- [BLS] M.Bożejko, M.Leinert, R.Speicher Convolution and limit theorems for conditionally free random variables, *Pacific J.Math.* **175** (1996), 357-388.
- [BW] M.Bożejko, J.Wysoczański, Remarks on t-transformations of measures and convolutions, *Ann. I.H.Poincare* **37** (2001), 737-761.
- [FL] U.Franz, R.Lenczewski, Limit theorems for the hierarchy of freeness, *Probab. Math. Stat.* **19** (1999), 23-41.
- [L1] R.Lenczewski, Unification of independence in quantum probability, *Inf. Dim. Anal. Quant. Prob. Rel. Topics*, **1** (1998), 383-405.
- [L2] R.Lenczewski, Reduction of free independence to tensor independence, *Inf. Dim. Anal. Quant. Prob. Rel. Topics.* **7** (2004), 337-360.
- [Lu] Y. G. Lu, On the interacting Fock space and the deformed Wigner law", *Nagoya Math. J.* **145** (1997), 1-28.
- [M1] N.Muraki, Noncommutative Brownian motion in monotone Fock space, *Commun. Math. Phys.* **140** (1997), 557-570.
- [M2] N.Muraki, Monotonic independence, monotonic central limit theorem and monotonic law of small numbers, *Inf. Dim. Anal. Quant. Prob. Rel. Topics.* **4** (2001), 39-58.
- [S] R. Speicher, A new example of "independence" and "white noise", *Probab. Th. Rel. Fields* **84** (1990) , 141-159.
- [V] D.Voiculescu, Symmetries of some reduced free product C^* -algebras, Operator Algebras and their Connections with Topology and Ergodic Theory, *Lecture Notes in Math.* 1132, Springer, Berlin, 1985, 556-588.
- [VDN] D.Voiculescu, K.Dykema, A.Nica, *Free Random Variables*, AMS, 1992.